

JointHealth™

Medications Guide March 2015

Arthritis Consumer Experts' (ACE) members/patients-at-large consistently request evidence-based information to enable them to have a more meaningful conversation with their rheumatologist and pharmacist around therapy of choice and route of administration.

In response, ACE produces an annual Medications Guide. The medications listed below are the most commonly prescribed by Canadian rheumatologists and arthritis specialists to treat osteoarthritis (OA), rheumatoid arthritis (RA), axial spondyloarthritis (which includes ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis), juvenile idiopathic arthritis (JIA), psoriatic arthritis (PsA), osteoporosis, systemic lupus erythematosus (SLE), and vasculitis.

The information in this JointHealth™ Medications Guide is not intended to suggest a course of treatment. It is for information only. Always speak to your doctor before starting or stopping a medication.

Important notes

acetaminophen
= Tylenol®

Generic names of medications are shown in lower case and the first letter of brand names is capitalized.



Dosages are not provided as they vary based on disease severity and individual need.



Only the most common and the most serious possible side effects are highlighted.



General information on public drug plan coverage is provided. For private drug plan coverage, please contact your benefits provider for information.

Medication	Symptoms and diseases commonly used to treat	Most common and most serious side effects	Public drug plan coverage
disease-modifying anti-rheumatic drugs (DMARDs)			
azathioprine (Imuran®) – pill	Inflammation and pain caused by, SLE and vasculitis. Effective at treating the underlying disease process in, SLE, vasculitis.	Most common include: Stomach upset. Most serious include: Increased risk of infection, low blood counts (bone marrow suppression), mouth ulcers, liver and pancreas toxicity. Blood tests must be done regularly to check blood counts and liver tests. Serious drug interactions can occur with allopurinol. When the azathioprine dose is not adjusted with the allopurinol dose, death can occur.	All plans offer coverage
hydroxychloroquine sulfate (Plaquenil®) – pill Should see an improvement in 3 to 6 months, and improvement can continue up to a year	Inflammation and pain caused by RA, PsA, SLE, OA.	Most common include: Stomach upset, cramps and diarrhea. Long-term use can cause skin pigmentation changes. Itchy skin rash (usually within a couple of weeks); many types of skin rashes. Rare: Irritability, nightmares, headaches, blurred vision, vision halo. Most serious include: Rare retinal (eye) toxicity in 1 out of 5000 – stay under 6mg/kg/day. The dose is based on lean body weight. Yearly eye exams are recommended. Low blood counts. Rare nerve/muscle dysfunction. Tinnitus.	All plans offer coverage
leflunomide (Arava®) – pill Should see an improvement in 1 to 2 months	Inflammation and pain caused by RA. Effective at treating the underlying disease process in RA, PsA.	Most common include: Stomach upset, diarrhea, increased risk of infection, high blood pressure, headache, skin rash. Most serious include: Liver toxicity and severe liver damage leading to death, severe infection, low blood counts, nerve damage, teratogenic (can cause malformations of developing fetus). Blood tests must be done regularly to check blood counts and liver tests. Leflunomide stays in the body for many months. If there is a serious side effect, or there is a plan to become pregnant, this drug should be “washed out” of the system by using a course of cholestyramine.	All plans offer coverage
methotrexate (Rheumatrex®) – pill or 1 injection weekly Benefits should be seen within 1 to 2 months; maximum benefit in 6 months	Inflammation and pain caused by RA, AS, PsA, SLE. Effective at treating the underlying disease process in RA, AS, SLE, PsA – peripheral arthritis only.	Most common include: Mouth ulcers, stomach upset, nausea, diarrhea, headache, fatigue, mood symptoms. Some patients may experience worsening of RA nodules. Most serious include: Liver toxicity, lung toxicity, low blood counts, increased risk of infection, hair loss, teratogenic (can cause malformations of developing fetus). Blood tests must be done regularly to check blood counts and liver tests.	All plans offer coverage
sulfasalazine (Azulfidine®) – pill Benefits should be seen in 1 to 2 months	Inflammation and pain caused by RA, AS, PsA. Effective at treating the underlying disease process in RA, AS – peripheral arthritis only.	Most common include: Nausea, stomach upset, diarrhea, abdominal pain, skin rash. Most serious include: Liver toxicity, drop in blood counts, temporary drop in sperm counts. Can cause kidney stones. Blood tests must be done regularly to check blood counts and liver tests.	All plans offer coverage

Medication	Symptoms and diseases commonly used to treat	Most common and most serious side effects	Public drug plan coverage
biologic response modifiers (biologics) – not to be used in combination with one another			
abatacept (Orencia®) – intravenous, at week 0, 2 and 4, and then once every 4 weeks or one injection per week.	Inflammation, pain, joint damage caused by RA, JIA. Highly effective at treating symptoms and underlying disease process in RA, JIA.	Most common include: Infusion reactions can occur and are usually mild and self limiting. Also, headache, runny nose, muscle or joint pain, sore throat, nausea, dizziness, heartburn. Most serious include: Increased risk of serious infections, possible increased risk of lymphoma, reactivation of hepatitis B, reactivation of tuberculosis (TB).	See Arthritis Medications Report Card
adalimumab (Humira®) – one injection every 2 weeks	Inflammation, pain, joint damage caused by RA, AS, PsA, JIA. Highly effective at treating symptoms and underlying disease process in RA, AS, PsA, JIA.	Most common include: Headache, skin rash, injection site reactions, rash, increased risk of minor infections. Most serious include: Low blood counts, increased risk of serious infection,, multiple sclerosis-like symptoms, possible increased risk of lymphoma, reactivation of hepatitis B, reactivation of tuberculosis (TB). Should not be used in people with severe or uncontrolled heart failure.	See Arthritis Medications Report Card
anakinra (Kineret®) – one injection every day	Inflammation and pain caused by RA, adult Still's disease.	Most common include: Injection site reactions, skin rash, headache, nausea, abdominal pain, increased risk of minor infections. Most serious include: Increased risk of serious infection, reactivation of hepatitis B, reactivation of tuberculosis (TB).	See Arthritis Medications Report Card
belimumab (Benlysta®) – intravenous infusion, every 2 weeks for the first 3 doses, then once every 4 weeks	Effective at treating symptoms and underlying disease process in systemic lupus erythematosus (SLE). Currently approved to treat skin and joint manifestations. Not indicated for renal or central nervous system (CNS) lupus.	Most common include: Nausea, diarrhea, fever, stuffy or runny nose, cough (bronchitis), trouble sleeping, leg or arm pain, depression, headache (migraine), sore throat, urinary tract infection, decreased white blood cell count (leukopenia), vomiting, stomach pain. Most serious include: Possible cancer; allergic and infusion reactions; serious reactions may occur on the day or day after receiving dose, and may cause death; infections; heart problems; mental health problems, including suicide, reactivation of hepatitis B, reactivation of tuberculosis (TB).	See Arthritis Medications Report Card
certolizumab pegol (Cimzia®) – one injection every 2 weeks	Inflammation, pain, joint damage caused by RA, AS, PSA. Highly effective at treating symptoms and underlying disease process in RA, AS, PSA.	Most common include: Upper respiratory tract infections, rash, urinary tract infections, lower respiratory tract and lung infections. Most serious include: Infections including malignancies including possible increased risk of lymphoma, reactivation of hepatitis B, reactivation of tuberculosis (TB).	See Arthritis Medications Report Card
denosumab (Prolia®) – injection, 2 per year	Osteoporosis in postmenopausal women who have a high risk of bone fractures.	Most common include: Back pain, pain in arms and legs, high cholesterol, muscle pain, and bladder infection. Note: These events were also very common in the placebo group. Most serious, but rare, include: Infections in skin, lower stomach area (abdomen), bladder, or ear; inflammation of inner lining of heart (endocarditis) due to infection; osteonecrosis of the jaw (very rare); lowered calcium levels in blood (hypocalcemia), reactivation of hepatitis B, reactivation of tuberculosis (TB).	See Arthritis Medications Report Card

biologics continued

Medication	Symptoms and diseases commonly used to treat	Most common and most serious side effects	Public drug plan coverage
biologic response modifiers (biologics) – not to be used in combination with one another <i>continued</i>			
etanercept (Enbrel®) – one or two injections every week	Inflammation, pain, joint damage caused by RA, AS, JIA, PsA. Highly effective at treating symptoms and underlying disease process in RA, AS, JIA, PsA.	Most common include: Headache, skin rash, injection site reactions, rash, increased risk of minor infections, dizziness. Most serious include: Low blood counts, increased risk of serious infection, multiple sclerosis-like symptoms, possible increased risk of lymphoma, reactivation of hepatitis B, reactivation of tuberculosis (TB). Should not be used in people with severe or uncontrolled heart failure.	See Arthritis Medications Report Card
golimumab (Simponi®) – one injection every 4 weeks or for RA, intravenous infusion at weeks 0 and 4, then every 8 weeks	Inflammation, pain, joint damage caused by RA, AS, PsA. Highly effective at treating symptoms and underlying disease process in RA, AS, PsA.	Most common include: Upper respiratory tract infection, nausea, abnormal liver tests, redness at site of injection, high blood pressure, bronchitis, dizziness, sinus infection, flu, runny nose, fever, cold sores, numbness or tingling. Most serious include: Serious infection, increased risk of lymphoma, reactivation of TB, reactivation of hepatitis B, heart failure, nervous system problems, liver problems, blood problems.	See Arthritis Medications Report Card
infliximab (Remicade®) – intravenous infusion once every 8 weeks	Inflammation, pain, joint damage caused by RA, AS, PsA. Highly effective at treating symptoms and underlying disease process in RA, AS, PsA.	Most common include: Headache, skin rash, infusion reactions, rash, increased risk of minor infections. Most serious include: Low blood counts, increased risk of serious infection, multiple sclerosis-like symptoms, possible increased risk of lymphoma, reactivation of hepatitis B, reactivation of tuberculosis (TB). Should not be used in people with severe or uncontrolled heart failure.	See Arthritis Medications Report Card
rituximab (Rituxan®) – intravenous. For RA, the first two infusions are separated by 2 weeks, then usually re-infusion will occur every six months. For vasculitis, it is given weekly for 4 weeks. If or when another course is needed is not yet defined.	Inflammation, pain, joint damage caused by RA; used to treat ANCA vasculitis and can reduce or prevent organ damage.	Most common include: Infusion reactions are usually seen at first infusion, include flushing, sweating, chest pains. Infusion reactions are typically managed by slowing the rate of infusion and are much less frequent in subsequent infusions. Most serious include: Sore throat, fever, chills, or other signs of infection, unusual bruising or bleeding, severe pain in the stomach area, vision changes, unusual eye movements, loss of balance or coordination, confusion, disorientation, difficulty walking, risk of serious infection, reactivation of hepatitis B, reactivation of tuberculosis (TB). Higher risk of hepatitis B reactivation compared to other biologics.	See Arthritis Medications Report Card
tocilizumab (Actemra®) – intravenous infusion once every 4 weeks OR — subcutaneous injection every one to two weeks	Inflammation, pain, joint damage caused by RA and JIA. Highly effective at treating symptoms and underlying disease process in RA and JIA.	Most common include: Upper respiratory tract infection, nasopharyngitis, headache, hypertension, and increased ALT. Most serious include: Infections, in some cases fatal, gastrointestinal perforations, and hypersensitivity reactions including anaphylaxis, reactivation of hepatitis B, reactivation of tuberculosis (TB).	See Arthritis Medications Report Card
ustekinumab (Stelara®) – injection, at weeks 0 and 4, then every 12 weeks	Inflammation, pain, joint damage caused by PsA. Highly effective at treating symptoms and underlying disease process in PsA.	Most common include: Upper respiratory infections, headache, fatigue. Most serious include: Increased risk of infection, including reactivation of hepatitis B, reactivation of tuberculosis (TB); increased risk of certain types of cancer.	See Arthritis Medications Report Card

Medication	Symptoms and diseases commonly used to treat	Most common and most serious side effects	Public drug plan coverage
targeted small molecule medications (TSMMs)			
tofacitinib citrate (Xeljanz®) – pill, once morning and evening	Inflammation and pain caused by RA	Most common include: Nausea, indigestion, and diarrhea; headaches; upper respiratory tract infection (nasopharyngitis); may increase cholesterol levels. Most serious include: Infection, changes to red and white blood cell counts; may irritate the liver; may increase risk of bowel perforation; and may cause slight decrease in kidney function (increased creatinine).	See Arthritis Medications Report Card
subsequent entry biologics (SEBs)			
infliximab (Inflectra®) – intravenous infusion once every 8 weeks	Inflammation and pain caused by RA, AS, and PsA	Most common include: Abdominal pain, nausea, vomiting, and diarrhea; back pain, aching joints; rash, flushing; headaches; upper respiratory tract infections, such as sinusitis. Most serious, but rare, include: Infusion reactions; increased risk of infection, nervous system disorders' making congestive heart failure worse; and malignancy. Should not be used in people with severe or uncontrolled heart failure.	See Arthritis Medications Report Card
acetaminophen			
Examples – pill Tylenol®, Panadol®, Aspirin Free Anacin® Should see benefit within 30 minutes	Pain caused by RA, AS, PsA, SLE, osteoarthritis (OA) Does not treat the underlying disease process.	Most common include: Few common side effects. Rare side effects are rash, low blood counts, stomach upset. Most serious include: Sudden liver failure (large overdose) or chronic liver failure if used at higher than recommended doses, with alcohol, or with other liver-toxic drugs.	No
non-steroidal anti-inflammatory drugs (NSAIDs)			
Examples – pill diclofenac (Voltaren®) ibuprofen (Motrin®) indomethacin (Indocid®) meloxicam (Mobic®) naproxen (Naprosyn®) Full benefit within 2 weeks	Inflammation and pain caused by RA, AS, PsA, OA. Does not treat the underlying disease process.	Most common include: Stomach upset and dyspepsia. Most serious include: Peptic ulcer disease (1-4% / year), kidney toxicity, increased risk of cardiovascular disease, modest worsening of underlying high blood pressure, liver toxicity, asthma, low blood counts, increased risk of bleeding, skin rash. Peptic ulcer risk is reduced if a proton pump inhibitor (PPI) is co-administered. Periodic tests for liver and renal side effects are recommended. Note: Most NSAID side effects are “dose-dependent” so you should try to use the lowest effective dose “as needed”, rather than regularly. All NSAIDs currently available carry an increased risk of heart attack and stroke.	Most plans offer coverage
COX-2 NSAIDs			
celecoxib (Celebrex®) – pill Full benefit seen within 2 weeks	Inflammation and pain caused by RA, AS, PsA, OA. Does not treat the underlying disease process.	Most common include: Same as NSAIDs, except there may be less dyspepsia and stomach upset (often better tolerated). Most serious include: Same as NSAIDs except – 1. There is a reduced risk (about half the risk) of peptic ulcer disease and, – 2. There is a possible increased cardiovascular risk at the higher dose (200 mg twice daily). The patient who would benefit from Celebrex® over the other NSAIDs would be the patient with a higher risk of peptic ulcer disease (such as a previous ulcer) who also has a low cardiovascular risk. Periodic blood tests to check the liver and blood counts are recommended for patients who are on chronic doses of these medications.	A number of plans offer coverage, some only offer restricted coverage

Medication	Symptoms and diseases commonly used to treat	Most common and most serious side effects	Public drug plan coverage
steroids			
<p>glucocorticoids – can be given by mouth, by intravenous infusion (for life or organ-threatening disease), by intramuscular injection, by injection directly into a joint or tendon sheath (when there is local inflammation)</p> <p>cortisone dexamethasone hydrocortisone medroxyprogesterone methylprednisolone prednisone prednisolone</p> <p>Benefits should be seen within 24 hours</p>	<p>Inflammation caused by RA, AS, PsA, SLE, vasculitis. Sometimes given by injection to any joint (including OA).</p> <p>A good bridge therapy before DMARDs take full effect.</p> <p>It is unlikely that glucocorticoids will slow down the joint damage caused by these diseases, but glucocorticoids are often used to treat life-threatening or organ-threatening complications of these diseases.</p> <p>For example: RA lung inflammation RA eye disease RA vasculitis.</p>	<p>Side effects are usually dose and time dependent. They rarely occur with single injections or short course, but are very frequent and sometimes irreversible with higher doses or long courses.</p> <p>Short term side effects include: Sleep disturbance, mood swings or even psychosis, blurred vision.</p> <p>The side effects listed below are generally seen with long-term use (at least a couple of months).</p> <p>Most common include: Stomach upset, thin skin, easy bruising, central weight gain, facial fullness (moon face) buffalo hump, increased hair growth, acne, thin extremities with muscle wasting and weakness, glaucoma, cataracts, increased cardiovascular risks, high cholesterol, high blood pressure, mood swings, depression, osteoporosis and increased risk of fracture, increased risk of infection, worsening of diabetes in known diabetics, or induction of diabetes in people already prone to developing it.</p> <p>The risk of osteoporosis (thin bones that break easily) may be reduced by taking appropriate amounts of calcium, vitamin D and certain medications that build bone.</p> <p>Rare but serious: Psychosis, severe depression, stroke, heart attack, pancreatitis, peptic ulcer disease, and osteonecrosis (can occur over short-term use). This is due to the interruption of blood to the end of a long bone (hip, knee or shoulder typically). This may cause complete destruction of the joint and is irreversible, usually. Osteonecrosis risk higher in SLE.</p> <p>Adrenal crisis: Long-term use of glucocorticoids usually suppresses adrenal gland function (makes cortisol that our bodies need). Therefore suddenly stopping or rapidly reducing glucocorticoids can cause "cortisol deficiency". Symptoms include loss of appetite, nausea, vomiting, abdominal pain, weakness, fatigue confusion or coma. There may be problems with the blood electrolytes (sodium and potassium). Adrenal crisis can even occur in a person who is still on glucocorticoids. It can be precipitated by surgery, trauma or an infection.</p> <p>For this reason, people on long-term glucocorticoids should have a bracelet or necklace indicating that they are on "prednisone" for example. This way, emergency personnel will know what to look for and to provide appropriate glucocorticoid doses.</p>	All plans offer coverage
other medications			
pregabalin (Lyrica®) – capsule taken by mouth	Widespread muscle pain caused by fibromyalgia.	<p>Most common include: Dizziness, sleepiness, weight gain, blurred vision, dry mouth, swelling of hands and feet, trouble concentrating.</p> <p>Most serious include: Serious allergic reactions, suicidal thoughts or actions, muscle problems, problems with eyesight, feeling "high".</p>	Most provinces do not yet list but resubmissions are ongoing
duloxetine (Cymbalta®)	Management of pain associated with fibromyalgia and for the chronic pain associated with OA of the knee.	<p>Most common include: Nausea, dizziness, fatigue, drowsiness, muscle weakness, constipation, dry mouth, diarrhea, abdominal pain, insomnia, decreased appetite, weight gain, and erectile dysfunction</p> <p>Most serious include: Serotonin syndrome and neuroleptic malignant syndrome (cause brain, muscle digestive system, and autonomic nervous system changes), liver disorder, Stevens-Johnson syndrome and erythema multiforme (serious skin reactions).</p> <p>Rare side effects include gastrointestinal bleeding, feeling of restlessness, glaucoma, mania, severe allergic reaction, weight gain, and low sodium level in blood.</p>	Most plans offer coverage.



Arthritis Consumer Experts™

#200A 1228 Hamilton Street, Vancouver BC V6B 6L2
t: 604.974.1366

e: feedback@jointhealth.org
www.jointhealth.org

Disclaimer

The material contained in this or any other ACE publication is provided for general information only. It should not be relied on to suggest a course of treatment for a particular individual or as a substitute for consultation with qualified health professionals who are familiar with your individual medical needs. If you have any healthcare related questions or concerns, you should contact your physician. Never disregard medical advice or delay in seeking it because of something you have read in any ACE publication.